

EXCEPTIONAL SEQUENCES OVER PATH ALGEBRAS OF TYPE A_n AND NON-CROSSING SPANNING TREES

TOKUJI ARAYA

ABSTRACT. Exceptional sequences are fundamental to investigate the derived categories of finite dimensional algebras. The aim of this note is to classify all the complete exceptional sequences over the path algebra of a Dynkin quiver of type A_n in terms of non-crossing spanning trees.

1. INTRODUCTION

The concept of exceptional sequences was introduced by Gorodentsev and Rudakov [10] to study exceptional vector bundles on $\mathbb{P}^2$. Exceptional sequences are certain variation of tilting objects and useful to study the structure of derived categories. They have effectively been applied to study the derived categories of algebraic varieties by many authors (e.g. [4], [17]). Also exceptional sequences were studied for the path algebras of acyclic quivers by Crawley-Boevey [6], for the weighted projective lines by Meltzer [15] and for Cohen-Macaulay modules over the one dimensional graded Gorenstein rings with a simple singularity by the author [1]. For example, the transitivity of the braid group action was shown for many cases. But their concrete description of the exceptional sequences does not seem to be studied yet even for the path algebras of Dynkin quivers.

In this paper, we give a combinatorial description of the exceptional sequences in the derived category of the path algebra Λ of type A_n . Let us introduce the notation which is necessary to state our results. Let $\epsilon = (E_1, E_2, \dots, E_n)$ and $\epsilon' = (E'_1, E'_2, \dots, E'_n)$ be complete exceptional sequences in $\mathcal{D}^b(\text{mod } \Lambda)$ (see 2.1 for the definition), we write $\epsilon \sim \epsilon'$ if there exists a permutation σ and integers $l_1, l_2, \dots, l_n$ such that $E_i \cong \Sigma^{l_i} E'_{\sigma(i)}$ for every i . Here, Σ is a shift functor in $\mathcal{D}^b(\text{mod } \Lambda)$. We denote by $\mathcal{E}$ the set of complete exceptional sequences in $\mathcal{D}^b(\text{mod } \Lambda)$, and we put $\overline{\mathcal{E}} = \mathcal{E} / \sim$.

Theorem 1.1. *There exists a bijection between $\overline{\mathcal{E}}$ and the set of non-crossing spanning trees on the circle with $(n+1)$ -points (defined in 2.3).*

We will explicitly give the bijection in terms of the Auslander-Reiten quivers. The proof of Theorem 1.1 depends on a traditional technique for combinatorics on Auslander-Reiten quivers introduced by Gabriel [9] and extensively studied by Riedtmann [17] and Wiedemann [19] for classification of the representation-finite selfinjective algebras and Gorenstein orders. Recently, this kind of combinatorics appeared also in cluster tilting theory by Caldero-Chapoton-Schiffler [5] and Iyama [14].

2000 *Mathematics Subject Classification.* 16G70, 05C05.

Key words and phrases. exceptional sequence, Dynkin quiver, Auslander-Reiten quiver, non-crossing spanning tree.

It is known the number of non-crossing spanning trees with $(n + 1)$ -points by Dulucq and Penaud [7] (see also [8]). Therefore we get following.

Corollary 1.2. *The cardinarity of $\overline{\mathcal{E}}$ is equal to $\frac{1}{2n+1} \binom{3n}{n}$.*

On the other hand, Seidel [18] determines that the number of exceptional sequences (in mod Λ) is equal to $(n + 1)^{n-1}$, but a concrete description of the exceptional sequences is not given there.

Also we will show in Section 5 that there is a nice interpretation of mutation of exceptional sequences in terms of non-crossing spanning trees.

2. PRELIMINARIES

Let Λ be the path algebra of a Dynkin quiver of type A_n over a field k . We denote by $\text{mod } \Lambda$ the category of finitely generated right Λ -modules and by $\mathcal{D}^b(\text{mod } \Lambda)$ the bounded derived category. We write Σ the shift functor of $\mathcal{D}^b(\text{mod } \Lambda)$.

Definition 2.1. An object $E \in \mathcal{D}^b(\text{mod } \Lambda)$ is called *exceptional* if $\text{Hom}(E, E) \cong k$ and $\text{Hom}(\Sigma^l E, E) = 0$ for all integers $l \neq 0$. A pair (E, F) of exceptional objects is called an *exceptional pair* if $\text{Hom}(\Sigma^l F, E) = 0$ for all integers l . A sequence $\epsilon = (E_1, E_2, \dots, E_r)$ of exceptional objects is called an *exceptional sequence* of length r if (E_i, E_j) is an exceptional pair for each $i < j$. An exceptional sequence ϵ is called *complete* if the length of ϵ is equal to the number n of simple Λ -modules.

We immediately get the following from the definition.


- Remark 2.2.** 1. The derived category $\mathcal{D}^b(\text{mod } \Lambda)$ is independent of the choice of orientation of A_n [12].
2. In general, exceptional objects are indecomposable. If Λ is the path algebra of a Dynkin quiver, the converse also holds. Thus, in our situation that Λ is the path algebra of A_n , an object is exceptional if and only if it is indecomposable.
 3. For any indecomposable object $X \in \mathcal{D}^b(\text{mod } \Lambda)$, there exists a unique integer l and a unique indecomposable Λ -module Y such that $X \cong \Sigma^l Y$ [12].
 4. Let $(E_1, E_2, \dots, E_n)$ be an exceptional sequence, then so is $(\Sigma^{l_1} E_1, \Sigma^{l_2} E_2, \dots, \Sigma^{l_n} E_n)$ for any integers $l_1, l_2, \dots, l_n$.
 5. Since $\text{gldim } \Lambda \leq 1$, we see that $E \in \text{mod } \Lambda$ is exceptional if $\text{Hom}(E, E) \cong k$ and $\text{Ext}_\Lambda^1(E, E) = 0$. Furthermore, we can see that (E, F) is an exceptional pair if $\text{Hom}(F, E) = \text{Ext}_\Lambda^1(F, E) = 0$ for all exceptional modules E and F .

By Remark 2.2.1, we only have to consider the case when the orientation is


$$\overset{1}{\bullet} \leftarrow \overset{2}{\bullet} \leftarrow \dots \leftarrow \overset{n}{\bullet}.$$

For any $(E_1, E_2, \dots, E_n) \in \overline{\mathcal{E}}$, by Remark 2.2.3 and 4, we may assume that all E_i are finitely generated Λ -modules.

Let $\Gamma = (\Gamma_0, \Gamma_1)$ be the Auslander-Reiten quiver (c.f. [3], [2]) of $\text{mod } \Lambda$. We identify the set Γ_0 of vertices in Γ with the isomorphism classes of indecomposable Λ -modules. For each integers $0 \leq i < j \leq n$, we write $X_{i,j}$ the indecomposable Λ -module whose representation is given by $(\overset{1}{0} \leftarrow \dots \leftarrow \overset{i}{0} \leftarrow \overset{i+1}{k} \leftarrow \dots \leftarrow \overset{j}{k} \leftarrow \overset{j+1}{0} \leftarrow \dots \leftarrow \overset{n}{0})$. Then it is known that $\Gamma_0 = \{X_{i,j} \mid 0 \leq i < j \leq n\}$ and Γ is the following form:



Now we consider the circle with $n + 1$ points labeled $0, 1, 2, \dots, n$ counter clockwise on it. We write $c(i, j)(= c(j, i))$ the chord between the points i and j . We denote by C_{n+1} the set of chords on the circle.




Definition 2.3. A set T consisting of n chords in C_{n+1} is called a *non-crossing spanning tree* if T satisfies the following conditions:

- (i) the chords in T make a tree,
- (ii) the chords in T meet only at their endpoints.


We remark that each vertex is then contained in at least one chord.

Here we give some examples of non-crossing spanning tree and not non-crossing spanning trees on the circle with 6 points.


Example 2.4. 1. The following diagram is a non-crossing spanning tree.



2. The following diagram is not a non-crossing spanning tree because it has a cycle.



3. The following diagram is not a non-crossing spanning tree because $c(0, 2)$ meets $c(1, 3)$ in the interior of the circle.



Since $\Gamma_0 = \{X_{i,j} | 0 \leq i < j \leq n\}$ and $C_{n+1} = \{c(i,j) | 0 \leq i < j \leq n\}$, there is a bijection

$$\Phi : \Gamma_0 \rightarrow C_{n+1} \quad (X_{i,j} \mapsto c(i,j)).$$

Now we can state Theorem 1.1 in the following more explicit form.

Theorem 2.5. *A bijection from $\overline{\mathcal{E}}$ to the set of non-crossing spanning trees on the circle with $(n+1)$ -points is given by*

$$\epsilon = (E_1, E_2, \dots, E_n) \mapsto \Phi(\epsilon) := \{\Phi(E_i) \mid 1 \leq i \leq n\}.$$

We shall give a proof of Theorem 2.5 in the next section.

3. PROOF OF THEOREM 2.5

For $X \in \Gamma_0$, we consider the following four subsets of Γ_0 .

$$\mathcal{H}_+^0(X) = \{Y \in \Gamma_0 \mid \text{Hom}(X, Y) \neq 0\},$$

$$\mathcal{H}_-^0(X) = \{Y \in \Gamma_0 \mid \text{Hom}(Y, X) \neq 0\},$$

$$\mathcal{H}_+^1(X) = \{Y \in \Gamma_0 \mid \text{Ext}_\Lambda^1(X, Y) \neq 0\},$$


$$\mathcal{H}_-^1(X) = \{Y \in \Gamma_0 \mid \text{Ext}_\Lambda^1(Y, X) \neq 0\},$$

and we put $\mathcal{H}_\pm(X) = \mathcal{H}_\pm^0(X) \cup \mathcal{H}_\pm^1(X)$.

Lemma 3.1. For $X = X_{i,j}$, the above four sets are given by the following forms in Γ .

$$\begin{aligned}\mathcal{H}_+^0(X_{i,j}) &= \{X_{s,t} \mid i \leq s \leq j-1, j \leq t \leq n\}, \\ \mathcal{H}_-^0(X_{i,j}) &= \{X_{s,t} \mid 0 \leq s \leq i, i+1 \leq t \leq j\}, \\ \mathcal{H}_+^1(X_{i,j}) &= \{X_{s,t} \mid 0 \leq s \leq i-1, i \leq t \leq j-1\}, \\ \mathcal{H}_-^1(X_{i,j}) &= \{X_{s,t} \mid i+1 \leq s \leq j, j+1 \leq t \leq n\}.\end{aligned}$$

Proof. For $X, Y \in \Gamma_0$, the dimension of the k -vector space $\text{Hom}(X, Y)$ can be calculated by using Auslander-Reiten sequences. Indeed, if there is no path from X to Y in Γ , then we have $\text{Hom}(X, Y) = 0$ (c.f. [3], [2]). If $0 \rightarrow \tau Y \rightarrow \bigoplus_{i=1}^r Z_i \rightarrow Y \rightarrow 0$ is an Auslander-Reiten sequence, then the equality $\dim_k \text{Hom}(X, Y) + \dim_k \text{Hom}(X, \tau Y) = \sum_{i=1}^r \dim_k \text{Hom}(X, Z_i)$ holds. Thus, we get $\dim_k \text{Hom}(X, Y) = 1$ if $\text{Hom}(X, Y) \neq 0$, and we can draw the areas $\mathcal{H}_\pm^0(X_{i,j})$ in Γ as follows.



Using the Auslander-Reiten duality $\text{Ext}_\Lambda^1(X, Y) \cong \text{D Hom}(Y, \tau X) (\cong \text{D Hom}(\tau^- Y, X))$ where $\text{D}(-)$ is the k -dual functor $\text{Hom}_k(-, k)$ (c.f. [3], [2]), we get $\mathcal{H}_+^1(X_{i,j}) = \mathcal{H}_-^0(\tau X_{i,j}) = \mathcal{H}_-^0(X_{i-1,j-1})$ and $\mathcal{H}_-^1(X_{i,j}) = \mathcal{H}_+^0(\tau^- X_{i,j}) = \mathcal{H}_+^0(X_{i+1,j+1})$.

□

For the points i and j on the circle, we can consider the distance $d(i, j)$ from i to j obtained by a map $d : \{0, 1, \dots, n\} \times \{0, 1, \dots, n\} \rightarrow \mathbb{N}$ which is given by

$$d(i, j) = \begin{cases} j - i & (j \geq i) \\ j + n + 1 - i & (j < i). \end{cases}$$

The following lemma is a key to prove Theorem 2.5.

Lemma 3.2. Let $X = X_{i,j}, X' = X_{i',j'} \in \Gamma_0$.

1. Both (X, X') and (X', X) are exceptional pairs if and only if $\Phi(X)$ does not meet $\Phi(X')$.
2. Neither (X, X') nor (X', X) is an exceptional pair if and only if $\Phi(X)$ meets $\Phi(X')$ in the interior of the circle.

3. (X, X') is an exceptional pair but (X', X) is not an exceptional pair if and only if one of the following conditions holds:

- (i) $i = i'$ and $d(i, j) < d(i', j')$,
- (ii) $i = j'$ and $d(i, j) < d(j', i')$,
- (iii) $j = i'$ and $d(j, i) < d(i', j')$,
- (iv) $j = j'$ and $d(j, i) < d(j', i')$.

Proof. From the definition of an exceptional pair, we see the following.

1. Both (X, X') and (X', X) are exceptional pairs if and only if $X' \in \Gamma_0 \setminus (\mathcal{H}_+(X) \cup \mathcal{H}_-(X))$.
2. Neither (X, X') nor (X', X) is an exceptional pair if and only if $X' \in \mathcal{H}_+(X) \cap \mathcal{H}_-(X)$.
3. (X, X') is an exceptional pair but (X', X) is not an exceptional pair if and only if $X' \in \mathcal{H}_+(X) \setminus \mathcal{H}_-(X)$.


We can obtain the desired conclusion by using Lemma 3.1. □

Now we can prove Theorem 2.5.


Proof of Theorem 2.5. Let $\epsilon = (E_1, E_2, \dots, E_n)$ be a complete exceptional sequence. We shall prove that $\Phi(\epsilon) = \{\Phi(E_1), \Phi(E_2), \dots, \Phi(E_n)\}$ is a non-crossing spanning tree. Since $\Phi(E_i)$ does not meet $\Phi(E_j)$ in the interior of the circle for all $i \neq j$, it is enough to show that $\Phi(\epsilon)$ does not have cycles. To show this, we assume that $\Phi(E_{l_1}), \Phi(E_{l_2}), \dots, \Phi(E_{l_r})$ ($l_1 < l_2 < \dots < l_r$) make cycle. Moreover, we may assume that $\Phi(E_{l_j})$ meets $\Phi(E_{l_{j+1}})$ at the point i_j for each $j = 1, 2, \dots, r$ with $E_{r+1} = E_1$ and that $d(i_j, i_{j+1}) > d(i_j, i_{j-1})$ with $i_0 = i_r$ and $i_{r+1} = i_1$. In particular, $\Phi(E_r)$ meets $\Phi(E_1)$ at i_r and $d(i_r, i_1) > d(i_r, i_{r-1})$. This contradicts that (E_1, E_r) is an exceptional pair and Lemma 3.2 (3). Therefore $\Phi(\epsilon)$ does not have cycles and it is a non-crossing spanning tree.

Let $T = \{c_1, c_2, \dots, c_n\}$ be a non-crossing spanning tree. Since c_i does not meet c_j in the interior of the circle, either $(\Phi^{-1}(c_i), \Phi^{-1}(c_j))$ or $(\Phi^{-1}(c_j), \Phi^{-1}(c_i))$ is an exceptional pair for all $i \neq j$. We remark that there exists $c_{i_1} \in T$ such that $(\Phi^{-1}(c_{i_1}), \Phi^{-1}(c))$ is an exceptional pair for all $c \in T \setminus \{c_{i_1}\}$. Indeed, if such c_{i_1} does not exist, then T must have a cycle. We put $E_1 = \Phi^{-1}(c_{i_1})$. We choose c_{i_j} such that $(\Phi^{-1}(c_{i_j}), \Phi^{-1}(c))$ is an exceptional pair for all $c \in T \setminus \{c_{i_1}, c_{i_2}, \dots, c_{i_j}\}$ inductively and put $E_j = \Phi^{-1}(c_{i_j})$. Then $(E_1, E_2, \dots, E_n)$ is a complete exceptional sequence. □

Example 3.3. If $n = 3$, the following quiver is the Auslander-Reiten quiver of $\text{mod } \Lambda$.




In this case, there are 16 complete exceptional sequences and 12 non-crossing spanning trees. The following are the complete exceptional sequences and the corresponding non-crossing spanning trees.



4. ACTION OF THE CYCLIC GROUP

Let $\mathbb{Z}/(n+1)\mathbb{Z} = \langle \sigma \rangle$ be the cyclic group of order $n+1$. The group $\mathbb{Z}/(n+1)\mathbb{Z}$ acts on C_{n+1} by

$$\sigma(c(i, j)) = \begin{cases} c(i-1, j-1) & (i \neq 0), \\ c(j-1, n) & (i = 0). \end{cases}$$



The group $\mathbb{Z}/(n+1)\mathbb{Z}$ also acts on Γ_0 via Φ and its action is given by

$$\sigma(X_{i,j}) = \begin{cases} X_{i-1,j-1} & (i \neq 0) \\ X_{j-1,n} & (i = 0) \end{cases} \quad \text{for } X_{i,j} \in \Gamma_0$$

Thus, we see

$$\sigma(X) = \begin{cases} \tau X & \text{if } X \text{ is not projective,} \\ \nu X & \text{if } X \text{ is projective,} \end{cases}$$

where ν is the Nakayama functor.

We set $\sigma(\epsilon) = (\sigma(E_1), \sigma(E_2), \dots, \sigma(E_n))$ for $\epsilon = (E_1, E_2, \dots, E_n) \in \mathcal{E}$. In general, we must check that $\sigma(\epsilon)$ is a complete exceptional sequence by using the Auslander-Reiten duality. But this is immediate from Theorem 2.5 since $\Phi(\sigma(\epsilon))$ is clearly a non-crossing spanning tree.

5. MUTATION OF EXCEPTIONAL SEQUENCES

Let (E, F) be an exceptional pair (in $\mathcal{D}^b(\text{mod } \Lambda)$). The *left mutation* $\mathcal{L}_E F$ of F by E is given by a distinguished triangle

$$\bigoplus_{l \in \mathbb{Z}} \text{Hom}(\Sigma^l E, F) \otimes_k \Sigma^l E \xrightarrow{\text{can}} F \rightarrow \mathcal{L}_E F,$$

where *can* denote the canonical morphism. Dually, the *right mutation* $\mathcal{R}_F E$ of E by F is given by a distinguished triangle

$$\mathcal{R}_F E \rightarrow E \xrightarrow{\text{can}} \bigoplus_{l \in \mathbb{Z}} \text{D Hom}(E, \Sigma^l F) \otimes_k \Sigma^l F.$$

We denote by $B_n = \langle \sigma_1, \sigma_2, \dots, \sigma_{n-1} \mid \sigma_i \sigma_j = \sigma_j \sigma_i \ (j - i \geq 2), \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \ (i = 1, 2, \dots, n-2) \rangle$ the braid group on n -strings and by $\mathbb{Z}^n = \bigoplus_{i=1}^n \mathbf{e}_i \mathbb{Z}$ a free abelian group of rank n . It is known that $G := B_n \ltimes \mathbb{Z}^n$ acts on $\mathcal{E}$ by

$$\begin{aligned} \sigma_i(\epsilon) &= (E_1, E_2, \dots, E_{i-1}, \mathcal{L}_{E_i} E_{i+1}, E_i, E_{i+2}, \dots, E_n), \\ \mathbf{e}_i(\epsilon) &= (E_1, E_2, \dots, E_{i-1}, \Sigma E_i, E_{i+1}, \dots, E_n). \end{aligned}$$

Crawley-Boevey [6] showed that this action is transitive, namely, for any $\epsilon, \epsilon' \in \mathcal{E}$, there exists $\rho \in G$ such that $\epsilon = \rho \epsilon'$.

In this section, we consider a mutation of chords which corresponds to exceptional pairs.

We extend $\Phi : \Gamma_0 = \{\text{indecomposable } \Lambda\text{-modules}\} \rightarrow C_{n+1}$ to $\Phi : \{\text{indecomposable objects in } \mathcal{D}^b(\text{mod } \Lambda)\} \rightarrow C_{n+1}$ by $\Phi(\Sigma X) = \Phi(X)$. By definition we have $\mathcal{L}_{\Sigma^i E} \Sigma^j F \cong \Sigma^j \mathcal{L}_E F$ and $\mathcal{R}_{\Sigma^j F} \Sigma^i E \cong \Sigma^i \mathcal{R}_F E$. Thus we obtain $\Phi(\mathcal{L}_{\Sigma^i E} \Sigma^j F) = \Phi(\mathcal{L}_E F)$ and $\Phi(\mathcal{R}_{\Sigma^j F} \Sigma^i E) = \Phi(\mathcal{R}_F E)$.


The following lemma is straightforward.

Lemma 5.1. *The following conditions are equivalent for an exceptional pair (E, F) :*

1. $\mathcal{L}_E F \cong F$;
2. $\mathcal{R}_F E \cong E$;
3. (F, E) is an exceptional pair;
4. $\bigoplus_{l \in \mathbb{Z}} \text{Hom}(\Sigma^l E, F) = 0$;
5. $\Phi(E)$ does not meet $\Phi(F)$.

From now on, we assume that (E, F) is an exceptional pair but (F, E) is not an exceptional pair. Note that $\Phi(E)$ meets $\Phi(F)$ at some point i by Lemma 3.2 (3).

Lemma 5.2. *Let (E, F) be an exceptional pair with $\Phi(E) = c(i, j)$ and $\Phi(F) = c(i, l)$. Then $\Phi(\mathcal{L}_E F) = \Phi(\mathcal{R}_F E) = c(j, l)$.*



Proof. There exist indecomposable Λ -modules X and Y , and integers a and b such that $E \cong \Sigma^a X$ and $F \cong \Sigma^b Y$. Since $\text{Hom}(\Sigma^l E, F) \cong \text{Hom}(\Sigma^l \Sigma^a X, \Sigma^b Y) \cong \text{Hom}(\Sigma^{l+a-b} X, Y)$ and

$$\text{Hom}(\Sigma^{l+a-b} X, Y) \cong \begin{cases} \text{Hom}(X, Y) & (l+a-b=0), \\ \text{Ext}_\Lambda^1(X, Y) & (l+a-b=-1), \\ 0 & \text{otherwise,} \end{cases}$$

we have $\sum_{l \in \mathbb{Z}} \dim_k \text{Hom}(\Sigma^l E, F) \leq 1$ by Lemma 3.1. The assumption implies $\sum_{l \in \mathbb{Z}} \dim_k \text{Hom}(\Sigma^l E, F) = 1$. We may assume that $\text{Hom}(E, F) \cong k$ and that f is a generator of $\text{Hom}(E, F)$. Then, $\mathcal{L}_E F$ and $\mathcal{R}_F E$ are given by the following triangles:

$$\begin{aligned} E &\xrightarrow{f} F \rightarrow \mathcal{L}_E F, \\ \mathcal{R}_F E &\rightarrow E \xrightarrow{f} F. \end{aligned}$$

Therefore we get $\mathcal{R}_F E \cong \Sigma^{-1} \mathcal{L}_E F$ and $\Phi(\mathcal{R}_F E) = \Phi(\mathcal{L}_E F)$.

To see $\Phi(\mathcal{L}_E F) = \Phi(\mathcal{R}_F E) = c(j, l)$, we assume E and F are Λ -modules.

Case 1. If $i < l < j$, then we have $E = X_{i,j}$ and $F = X_{i,l}$. Since $l < j$, we see $\text{Hom}(F, E) \neq 0$ by Lemma 3.1.

Case 2. If $l < j < i$, then we have $E = X_{j,i}$ and $F = X_{l,i}$. Since $l < j$, we see $\text{Hom}(F, E) \neq 0$ by Lemma 3.1.

Case 3. If $j < i < l$, then we have $E = X_{j,i}$ and $F = X_{i,l}$. Since $j < l-1$, we see $\text{Ext}_\Lambda^1(F, E) \neq 0$ by Lemma 3.1.

Thus these three cases contradict that (E, F) is an exceptional pair.

We remark that the representation of the indecomposable Λ -module $X_{s,t}$ is given by $(0 \leftarrow \dots \leftarrow \overset{s}{0} \leftarrow \overset{s+1}{k} \leftarrow \dots \leftarrow \overset{t}{k} \leftarrow \overset{t+1}{0} \leftarrow \dots \leftarrow \overset{n}{0})$.

Case 4. We assume $i < j < l$. Then we have $E = X_{i,j}$ and $F = X_{i,l}$. In this case, the generator of $\text{Hom}(E, F)$ is a monomorphism and $\mathcal{L}_E F$ is obtained by a short exact sequence $0 \rightarrow E \rightarrow F \rightarrow \mathcal{L}_E F \rightarrow 0$. Therefore each representations are given by

$$\begin{array}{cccccccccccccccc}
E & : & \cdots \leftarrow & 0 & \leftarrow & k^{i+1} & \leftarrow \cdots \leftarrow & k^j & \leftarrow & k^{j+1} & \leftarrow \cdots \leftarrow & 0^l & \leftarrow & 0^{l+1} & \leftarrow \cdots \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
F & : & \cdots \leftarrow & 0 & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & 0 & \leftarrow \cdots \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
\mathcal{L}_E F & : & \cdots \leftarrow & 0 & \leftarrow & 0 & \leftarrow \cdots \leftarrow & 0 & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & 0 & \leftarrow \cdots,
\end{array}$$

and we have $\mathcal{L}_E F = X_{j,l}$.

Case 5. We assume $j < l < i$. Then we have $E = X_{j,i}$ and $F = X_{l,i}$. In this case, the generator of $\text{Hom}(E, F)$ is an epimorphism and $\mathcal{R}_F E$ is obtained by a short exact sequence $0 \rightarrow \mathcal{R}_F E \rightarrow E \rightarrow F \rightarrow 0$. Therefore each representations are given by

$$\begin{array}{cccccccccccccccc}
\mathcal{R}_F E & : & \cdots \leftarrow & 0^j & \leftarrow & k^{j+1} & \leftarrow \cdots \leftarrow & k^l & \leftarrow & 0^{l+1} & \leftarrow \cdots \leftarrow & 0^i & \leftarrow & 0^{i+1} & \leftarrow \cdots \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
E & : & \cdots \leftarrow & 0 & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & 0 & \leftarrow \cdots \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
F & : & \cdots \leftarrow & 0 & \leftarrow & 0 & \leftarrow \cdots \leftarrow & 0 & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & 0 & \leftarrow \cdots,
\end{array}$$

and we have $\mathcal{R}_F E = X_{j,l}$.

Case 6. We assume $l < i < j$. Then we have $E = X_{i,j}$ and $F = X_{l,i}$. In this case, $\text{Ext}_\Lambda^1(E, F) \cong k$ and $\mathcal{L}_F E$ is obtained by a short exact sequence $0 \rightarrow F \rightarrow \mathcal{L}_F E \rightarrow E \rightarrow 0$. Therefore each representations are given by

$$\begin{array}{cccccccccccccccc}
F & : & \cdots \leftarrow & 0 & \leftarrow & k^{l+1} & \leftarrow \cdots \leftarrow & k^i & \leftarrow & 0^{i+1} & \leftarrow \cdots \leftarrow & 0^j & \leftarrow & 0^{j+1} & \leftarrow \cdots \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
\mathcal{L}_E F & : & \cdots \leftarrow & 0 & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & 0 & \leftarrow \cdots \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & \\
E & : & \cdots \leftarrow & 0 & \leftarrow & 0 & \leftarrow \cdots \leftarrow & 0 & \leftarrow & k & \leftarrow \cdots \leftarrow & k & \leftarrow & 0 & \leftarrow \cdots,
\end{array}$$

and we have $\mathcal{L}_E F = X_{l,j}$.

Thus we have $\Phi(\mathcal{L}_E F) = \Phi(\mathcal{R}_F E) = c(j, l)$.

□

Now we give a definition of mutation of chords which corresponds to exceptional pairs.

Definition 5.3. For $c, c' \in C_{n+1}$, we define the *left mutation* $\mathcal{L}_c c'$ of c' by c and the *right mutation* $\mathcal{R}_{c'} c$ of c by c' as follows:

1. If c does not meet c' , then we define $\mathcal{L}_c c' = c'$ and $\mathcal{R}_{c'} c = c$.
2. If $c = c(i, j)$ and $c' = c(i, l)$, then we define $\mathcal{L}_c c' = \mathcal{R}_{c'} c = c(j, l)$.

We can check the following Theorem by using Lemma 5.1 and Lemma 5.2.

Theorem 5.4. For an exceptional pair (E, F) , we have $\Phi(\mathcal{L}_E F) = \mathcal{L}_{\Phi(E)} \Phi(F)$ and $\Phi(\mathcal{R}_F E) = \mathcal{R}_{\Phi(F)} \Phi(E)$.

Acknowledgments. The author thanks Osamu Iyama and Frederic Chapoton for their comments and suggestions. He thanks Hugh Thomas for pointing out that this paper is closely related to [13] and [11]. He also thanks Ryo Takahashi for advice.

REFERENCES

- [1] T. Araya, *Exceptional sequences over graded Cohen-Macaulay rings*, Math. J. Okayama Univ. **41** (1999), pp 81-102
- [2] I. Assem, D. Simson, and A. Skowronski, *Elements of the representation theory of associative algebras Vol. 1. Techniques of representation theory*, London Mathematical Society Student Texts, 65. Cambridge University Press, Cambridge, 2006.
- [3] M. Auslander, I. Reiten and S. O. Smalø, *Representation theory of Artin algebras*, Cambridge Studies in Advanced Mathematics, 36. Cambridge University Press, Cambridge, 1995.
- [4] A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1183–1205, 1337; translation in Math. USSR-Izv. **35** (1990), no. 3, 519–541
- [5] P. Caldero, F. Chapoton and R. Schiffler, *Quivers with relations arising from clusters (A_n case)*, Trans. Amer. Math. Soc. **358** (2006), no. 3, 1347–1364.
- [6] E. Crawley-Boevey, *Exceptional sequences of representations of quivers*, Proceedings of the Sixth International Conference on Representations of Algebras (Ottawa, ON, 1992), 7, 117–124, Carleton-Ottawa Math. Lecture Note Ser., 14, Carleton Univ., Ottawa, ON, 1992.
- [7] S. Dulucq and J.-G. Penaud, *Cordes, arbres et permutations*, Discrete Math. **117** (1993), 89–105.
- [8] P. Flajolet and M. Noy, *Analytic combinatorics of non-crossing configurations*, Discrete Mathematics **204** (1999), 203–229
- [9] P. Gabriel, *Auslander-Reiten sequences and representation-finite algebras*, Representation theory, I (Proc. Workshop, Carleton Univ., Ottawa, Ont., 1979), pp. 1–71, Lecture Notes in Math., 831, Springer, Berlin, 1980.
- [10] A. L. Gorodentsev and A. N. Rudakov, *Exceptional vector bundles on projective spaces*, Duke. math. J. **54** (1987), 115–130.
- [11] I. Goulden and A. Yong, *Tree-like properties of cycle factorizations*, J. Combinatorial Theory (A) **98**, 2002, 106–117.
- [12] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*, London Mathematical Society Lecture Note Series, 119. Cambridge University Press, Cambridge, 1988
- [13] C. Ingalls and H. Thomas, *Noncrossing partitions and representations of quivers*, Preprint (2008), <http://arxiv.org/abs/math/0612219v4>.
- [14] O. Iyama, *Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories*, Adv. Math. **210** (2007), no. 1, 22–50.
- [15] H. Meltzer, *Exceptional vector bundles, tilting sheaves and tilting complexes for weighted projective lines*, Mem. Amer. Math. Soc. **171** (2004), no. 808
- [16] C. Riedtmann, *Representation-finite self-injective algebras of class A_n* , Representation theory, II (Proc. Second Internat. Conf., Carleton Univ., Ottawa, Ont., 1979), pp. 449–520, Lecture Notes in Math., 832, Springer, Berlin, 1980.
- [17] A. N. Rudakov, *Exceptional collections, mutations and helices*, Helices and vector bundles, 1–6, London Math. Soc. Lecture Note Ser., 148, Cambridge Univ. Press, Cambridge, 1990.
- [18] U. Seidel, *Exceptional sequences for quivers of Dynkin type*, Comm. Algebra **29** (3) (2001), 1373–1386.
- [19] A. Wiedemann, *Die Auslander-Reiten Köher der gitterendlichen Gorensteinordnungen*, Bayreuth. Math. Schr. No. **23** (1987), 1–134.

NARA UNIVERSITY OF EDUCATION, TAKABATAKE-CHO, NARA 630-8528, JAPAN
 E-mail address: araya@math.okayama-u.ac.jp